

# **AICHOR**

## Avicenna Journal of Care and Health in Operating Room

Avicenna J Care Health Oper Room. 2025;3(1):16-20. doi:10.34172/ajchor.84 https://ajchor.umsha.ac.ir



Original Article



# The Effect of Empathy Skill Training on Therapeutic Communication of Nursing Students in the Care of Hospitalized Children

Mohammad Nouri<sup>10</sup>, Neda Pashaei<sup>10</sup>, Monireh Rezaee Moradali<sup>20</sup>, Esfandyar Baljani<sup>1</sup>

- <sup>1</sup>Department of Nursing, Ur.C., Islamic Azad University, Urmia, Iran
- <sup>2</sup>Department of Midwifery, Ur.C., Islamic Azad University, Urmia, Iran

Article history: Received: February 24, 2025 Revised: March 27, 2025 Accepted: March 30, 2025 ePublished: April 29, 2025

\*Corresponding author: Esfandyar Baljani,

Email: Esfandyar.Baljani@iau.ir

#### Abstract

**Background:** Empathy is a crucial component of the nurse-patient relationship, serving as an essential clinical skill that strengthens the bond between nurses and patients and contributes to improved healthcare services. This study aimed to determine the impact of empathy skills training on the therapeutic communication of nursing students in caring for hospitalized children. **Methods:** This quasi-experimental study employed a single-group pretest-posttest design and was conducted on final-year nursing students at the Islamic Azad University of Urmia in 2023. The sample consisted of 40 nursing students selected through census sampling. At the beginning of the study, students completed a demographic information questionnaire and a therapeutic communication questionnaire developed by Han et al. Then, they received empathy skills training in groups over eight 90-minute sessions, held four times a week. The training was delivered through lectures using a video projector in small groups of six. After the intervention, students completed the questionnaires again. Data were analyzed using SPSS version 26.0. Descriptive statistics including mean (±SD) and frequency (percentage) were used to describe data, and paired *t*-test was used for statistical analysis.

**Results:** The mean overall therapeutic communication scores of nursing students were  $45.17 \pm 4.53$  before the intervention and  $47.97 \pm 4.43$  after the intervention. Based on the results of the paired *t*-test, there was a significant difference between the mean overall therapeutic communication scores before and after the intervention (t=-3.99, P<0.05).

**Conclusion:** Empathy skills training positively impacts the therapeutic communication of nursing students in caring for hospitalized children and can assist them in assessing the health problems of hospitalized children and providing better care.

Keywords: Empathy skills, Therapeutic communication, Nursing students, Children



**Please cite this article as follows:** Nouri M, Pashaei N, Rezaee Moradali M, Baljani E. The effect of empathy skill training on therapeutic communication of nursing students in the care of hospitalized children. Avicenna J Care Health Oper Room. 2025;3(1):16-20. doi:10.34172/ajchor.84

# Introduction

In clinical settings, the relationship between nurses and patients is key to ensuring that patients' needs are met (1,2). Empathy is defined as the ability to put oneself in another's position to better understand their feelings and experiences (3). It can reduce patients' defensiveness and encourage them to discuss their emotions and needs (4). It also shortens treatment duration, thereby reducing financial demands and resource needs (5). Nurses who care for patients empathetically can better understand patients' reactions to health issues and the sources and goals of these reactions (6,7). Empathy skills are highly valued among nursing students worldwide but require further

improvement, which is why nursing educators should pay more attention to fostering these skills in students (8).

A longitudinal study conducted by Ward et al on 214 nursing students found a significant decline in empathy from the beginning to the end of their training, particularly among students who had more clinical exposure to patients (9). Other studies on changes in empathy during education also indicate a decline in empathy as medical students progress through their studies (10), despite empathy being a fundamental aspect of all human interactions and a critical component of the nurse-patient relationship (11). A study conducted by Derksen et al in 2017 found that empathizing with patients



© 2025 The Author(s); Published by Hamadan University of Medical Sciences. This is an open-access article distributed under the terms of the Creative Commons Attribution License (https://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

calms and motivates them, increasing their willingness to cooperate, while a lack of empathy frustrates patients and discourages them from returning to healthcare providers (12). Nunes et al (13) and Ferri et al (14) also found that empathy among healthcare students declines over the course of their studies.

Communication is a key element between nurses, patients, and their families in healthcare. The primary goal of communication is to provide comfort and enhance feelings of trust and security in patients (15,16). Therapeutic communication is a planned conscious process aimed at improving patient outcomes (17). Nurses encourage patients to express their feelings, thoughts, and perceptions related to observable behaviors during communication (17,18). Implementing therapeutic communication methods by nurses not only leads to behavioral changes in themselves but also results in positive changes in patients' clinical conditions (19,20).

Many nursing students find communication skills challenging, which may lead to high levels of anxiety and hinder their ability to care for patients safely (21). Communication in pediatric wards is even more critical and challenging, as children at different developmental stages are more vulnerable and require better care (22). Therefore, healthcare workers must possess effective communication skills, particularly therapeutic communication skills, which include techniques to facilitate the comfortable expression of feelings and perceptions among patients and healthy individuals (23). For this reason, pediatric nursing education should include both theoretical and practical training in communication and therapeutic communication skills (24). Kudubes et al reported that the average therapeutic communication skills score was significantly low among nursing students who had completed pediatric internships (25). Similarly, a study by Loghmani et al found that nurses faced challenges in establishing appropriate communication with patients and their families due to overcrowding, time constraints, increased workload, financial issues, and workplace environment. Families expressed dissatisfaction with nurses' emotional responses, lack of reassurance, empathy, and failure to educate them (26).

A review of the literature indicates that studies on therapeutic communication with hospitalized children highlight positive outcomes such as patient participation in treatment, successful management of communication scenarios (27), and increased confidence and clinical performance among students (28). However, factors influencing and enhancing therapeutic communication have received less attention. Given the lack of studies on the impact of empathy skills training on the therapeutic communication of nursing students, this study aimed to determine the effectiveness of such training in the care of hospitalized children.

# **Materials and Methods**

This quasi-experimental study was conducted on final-

year nursing students at the Islamic Azad University of Urmia during the 2023-2024 academic year. Census sampling was used in the study, and the sample consisted of 40 final-year nursing students in a single group. Inclusion criteria included being a final-year nursing student in the internship phase and providing informed consent to participate in the study. The demographic questionnaire included age, gender, grade point average, marital status, and residency status. The therapeutic communication questionnaire by Han et al consisted of 15 items on a 4-point Likert scale. The tool comprised two domains: relationship building (9 items) and problemsolving (6 items) (29). The minimum and maximum scores were 15 and 60, respectively, with higher scores indicating better therapeutic communication. Cronbach's alpha was used to confirm the reliability of the tool, with an overall reliability of 0.89 (29).

To conduct the study, the researcher obtained a list of all final-year nursing students (in the internship phase) from the nursing faculty. Students completed the demographic and therapeutic communication questionnaires at the beginning of the study. Then, they were divided into groups of 6 and received eight 90-minute training sessions, held four times a week. Training was delivered through lectures using a video projector, and students' questions were addressed during the sessions. After the intervention, students completed the questionnaires again. The training content was based on the empathy skills training package from studies by Gholami et al (30) and Mohammadi Nesbat et al (31) (Table 1).

### Data Analysis

SPSS version 26.0 was used for data analysis in this study. Descriptive statistics were employed to describe demographic characteristics, including mean ( $\pm$  standard deviation) and frequency (percentage). The Shapiro-Wilk test was used to assess the normal distribution of quantitative data, and the paired t test was used to compare the therapeutic communication scores of students before and after the intervention.

#### **Results**

The results of the study showed that the frequency of nursing students participating in the study by gender included 29 females (72.5%) and 11 males (27.5%). The majority of them were in the age group of 22-27 years, and 36 (90%) were single. The demographic characteristics of the nursing students are presented in Table 2.

The results presented in Table 3 indicate that "empathy skills training" had a significant impact on improving the scores of therapeutic communications and all its dimensions, including relationship building, problemsolving, and overall therapeutic communication, among final-year nursing students.

As shown in Table 3, empathy skills training had a positive impact on the overall score of "therapeutic communication", particularly on the scores of the two

Table 1. Empathy Skills Training Package

Sessions	Topics	Session Contents
1	Sympathy	Definition of sympathy, benefits of sympathy, importance of sympathy, how to sympathize with others
2	Indifference	Definition of indifference, importance of indifference, difference between sympathy and indifference, role of indifference in social interactions
3	Self-awareness	Definition of self-awareness, importance and necessity of self-awareness skills, key points in developing self-awareness
4	Empathy skills	Introduction to empathy skills, stages of empathy development, empathy techniques, why empathy is the most effective life skill, effects of empathy
5	Acceptance of others	Becoming aware of judgments, methods to help individuals understand others (showing interest in others, accepting personalities, and respecting others)
6	Active listening	Listening effectively, reflecting emotions, five simple methods to cultivate students' sensitivity, reflecting content, understanding facial expressions and body language
7	Understanding others' perspectives	Understanding the other person's perspective, valuing others' personal viewpoints, accepting opposing perspectives, putting oneself in others' shoes.
8	Scenarios	Designing scenarios related to each session and discussing them with students

Table 2. Demographic Characteristics of the Students Participating in the Study

Variable	Category	Number (percent)		
Gender	Male	11 (27.5%)		
Gender	Female	29 (72.5%)		
Marital status	Single	36 (90%)		
Maritai status	Married	4 (10%)		
Davidson of the	Local	34 (85%)		
Residency status	Non-local	6 (15%)		
A	Mean	23.48		
Age	SD	1.32		
C	Mean	17.80		
Grade point average	SD	0.98		

**Table 3.** Comparison of Therapeutic Communication Scores of Final-Year Nursing Students Before and After the Intervention

Dimension	Time	Mean	SD	T Value	P value
Relationship	Before intervention	28.76	3.72	-3.93	< 0.001
building	After intervention	30.61	3.38		
D.II. I.	Before intervention	16.32	2.01	-2.25	0.030
Problem-solving	After intervention	17.27	2.06		
Overall therapeutic	Before intervention	45.17	4.53	-3.99	< 0.001
communication	After intervention	47.97	4.43		

domains of relationship building and problem-solving. The results of the paired t test indicated a significant difference between the mean therapeutic communication scores of nursing students before and after the empathy skills training (t=-3.99, P<0.001). The average overall score of therapeutic communication after intervention in this study was 47.97, indicating that nursing students had a higher level of therapeutic communication.

In the relationship-building domain, the mean scores before the intervention  $(28.76\pm3.72)$  compared to after the intervention  $(30.61\pm3.38)$  showed a significant difference (t=-3.93, P<0.001). This finding suggests that empathy skills training was effective in improving students' ability to establish initial relationships with patients.

Additionally, the results of the paired t test showed a

significant difference in the mean scores of students before and after the intervention in the problem-solving domain (t=-2.25, P<0.05). This indicates an improvement in students' ability to use communication skills to identify and resolve patients' problems.

#### Discussion

The results of the study demonstrated that empathy training was effective in enhancing the therapeutic communication skills of nursing students at the Islamic Azad University of Urmia. Given the significant improvement in overall therapeutic communication scores after the intervention, it can be concluded that improvement in one domain (e.g., relationship building) can contribute to improvement in other domains (e.g., problem-solving and overall therapeutic communication). This relationship highlights the synergistic effect of training on communication skills. In this study, the improvement in the problem-solving domain was less pronounced compared to the other domains. This difference may be due to the inherent complexity of problem-solving in the therapeutic communication process, which requires more practice and experience. The highest level of change in mean scores was observed in the relationship-building domain, indicating that the intervention was particularly effective in improving students' foundational skills in establishing relationships with patients. This skill, as the most basic aspect of therapeutic communication, holds significant importance (32).

Several studies have emphasized empathy as a crucial component of the communication process in medical sciences (33,34). Donovan et al taught therapeutic communication techniques through simulation (video clips) and scenario discussions in a study and found that these techniques increased nursing students' confidence, satisfaction, and perceived learning in the therapeutic communication process (35).

Managheb and Bagheri concluded that empathy skills training was effective in improving the performance of medical students, which is consistent with the results of the present study (36). Similarly, Spiro showed that

empathy is a teachable skill with a significant impact on medical students' performance, particularly during patient interviews (37). In the study titled "The Effectiveness of Empathy Skills Training on the Responsibility of Nursing Students", Mohammadi Nasab et al found that empathy skills training significantly increased the sense of responsibility among nursing students, which is in line with the present study (31).

The results of this study can be explained by the fact that empathy training in educational environments enhances the adaptability of nursing students and influences their ethical sensitivity, especially since the participants in this study were nursing students undergoing clinical training. These students had previously participated in professional ethics classes or workshops and had gained sufficient experience in interacting with patients during their years of study and patient care.

Regarding the practical implications of this study, it is worth noting that, compared to other skills, empathy, while somewhat dependent on personality, can be taught and learned, as demonstrated by this study and many similar studies. Empathy improves nurse-patient communication and accelerates the treatment process. Additionally, since the cognitive domain of students is influenced by education, nursing faculties can play a significant role in developing students' understanding of empathy and incorporating empathy skills as a specialized skill in therapeutic communication with hospitalized children. Therefore, empathy skills training should be integrated into the nursing curriculum, and workshops should be organized for nursing graduates to enhance patient satisfaction, reduce hospitalization time, improve health outcomes, and increase job satisfaction among nurses.

## Conclusion

The results of this study indicate that empathy has the potential to establish effective communication between patients and nursing students. Communication is the foundation of their care, so the communication between nursing students and patients helps patients understand their health conditions. Empathy skills training in the nursing student's curriculum can significantly improve the quality of therapeutic communication in the care of hospitalized children and have positive effects on the mental and physical health of patients. The results of this study emphasize that this skill, as an essential part of the nursing profession, should be continuously reinforced so that students can interact with patients in a more humane and effective manner. Ultimately, the results of this study suggest that empathy skills training can be considered a key factor in improving the quality of therapeutic communication and increasing patient satisfaction. However, it is necessary for these trainings to be structured and interactive, utilizing practical methods, so that nursing students can apply them in real-world settings, thereby enhancing the quality of care provided to patients.

#### Acknowledgments

The authors express their gratitude to the community staff for their invaluable contributions to the project execution and data collection. The authors also would like to thank nursing students for their participation in interviews.

#### **Authors' Contribution**

Conceptualization: Esfandyar Baljani.

Data curation: Mohammad Nouri, Neda Pashaei, Monireh Rezaee

Moradali.

Formal analysis: Mohammad Nouri.

Investigation: Mohammad Nouri, Neda Pashaei, Monireh Rezaee

Moradali.

**Methodology** Esfandyar Baljani. **Software**: Mohammad Nouri.

Supervision: Esfandyar Baljani, Monireh Rezaee Moradali.

Validation: Mohammad Nouri.

Writing-original draft: Esfandyar Baljani, Mohammad Nouri, Neda

Pashaei, Monireh Rezaee Moradali.

Writing-review & editing: Esfandyar Baljani, Monireh Rezaee Moradali.

#### **Competing Interests**

None to declare.

#### **Data Availability Statement**

The datasets used and/or analyzed during the current study are available from the corresponding author upon reasonable request.

#### **Ethical Approval**

This study was approved by the Ethics Committee of Urmia Islamic Azad University (IR.IAU.URMIA.REC.1403.050). Informed consent was obtained from all participants.

#### **Funding**

No particular grant from a public, private, or nonprofit organization was received for this research.

#### References

- Kahriman I, Nural N, Arslan U, Topbas M, Can G, Kasim S. The effect of empathy training on the empathic skills of nurses. Iran Red Crescent Med J. 2016;18(6):e24847. doi: 10.5812/ircmj.24847.
- Määttä SM. Closeness and distance in the nurse-patient relation. The relevance of Edith Stein's concept of empathy. Nurs Philos. 2006;7(1):3-10. doi: 10.1111/j.1466-769X.2006.00232.x.
- 3. Chen D, Lew R, Hershman W, Orlander J. A cross-sectional measurement of medical student empathy. J Gen Intern Med. 2007;22(10):1434-8. doi: 10.1007/s11606-007-0298-x.
- Mercer SW, Reynolds WJ. Empathy and quality of care. Br J Gen Pract. 2002;52 Suppl:S9-12.
- Williams B, Brown T, McKenna L, Boyle MJ, Palermo C, Nestel D, et al. Empathy levels among health professional students: a cross-sectional study at two universities in Australia. Adv Med Educ Pract. 2014;5:107-13. doi: 10.2147/amep.S57569.
- Reynolds WJ, Scott B. Do nurses and other professional helpers normally display much empathy? J Adv Nurs. 2000;31(1):226-34. doi: 10.1046/j.1365-2648.2000.01242.x.
- Haslam N. Empathy in patient care: antecedents, development, measurement, and outcomes by Mohammadreza Hojat. N Engl J Med. 2007;356(17):1792.
- Jia-Ru J, Yan-Xue Z, Wen-Nv H. Empathy ability of nursing students: a systematic review and meta-analysis. Medicine (Baltimore). 2022;101(32):e30017. doi: 10.1097/ md.000000000000030017.
- Ward J, Cody J, Schaal M, Hojat M. The empathy enigma: an empirical study of decline in empathy among undergraduate nursing students. J Prof Nurs. 2012;28(1):34-40. doi:

- 10.1016/j.profnurs.2011.10.007.
- Hojat M. Ten approaches for enhancing empathy in health and human services cultures. J Health Hum Serv Adm. 2009;31(4):412-50.
- 11. Cunico L, Sartori R, Marognolli O, Meneghini AM. Developing empathy in nursing students: a cohort longitudinal study. J Clin Nurs. 2012;21(13-14):2016-25. doi: 10.1111/j.1365-2702.2012.04105.x.
- 12. Derksen F, Olde Hartman TC, van Dijk A, Plouvier A, Bensing J, Lagro-Janssen A. Consequences of the presence and absence of empathy during consultations in primary care: a focus group study with patients. Patient Educ Couns. 2017;100(5):987-93. doi: 10.1016/j.pec.2016.12.003.
- Nunes P, Williams S, Sa B, Stevenson K. A study of empathy decline in students from five health disciplines during their first year of training. Int J Med Educ. 2011;2:12-7. doi: 10.5116/ ijme.4d47.ddb0.
- Ferri P, Guerra E, Marcheselli L, Cunico L, Di Lorenzo R. Empathy and burnout: an analytic cross-sectional study among nurses and nursing students. Acta Biomed. 2015;86 Suppl 2:104-15.
- Namdar H, Ebrahimi H, Yazdani H, Safaian A. Compliance in patients with schizophrenia. Iran Psychiatry Clinical Psychol. 2005;4(43):120. [Persian].
- 16. Shafakhah M, Zarshenas L, Sharif F, Sabet Sarvestani R. Evaluation of nursing students' communication abilities in clinical courses in hospitals. Glob J Health Sci. 2015;7(4):323-8. doi: 10.5539/gjhs.v7n4p323.
- 17. Mundakir M. Komunikasi Pelayanan Kesehatan. Indomedia Pustaka; 2016.
- Sarfika R, Maisa EA, Freska W. Buku Ajar Keperawatan Dasar
  Komunikasi Terapeutik Dalam Keperawatan. Padang: Andalas University Press; 2018.
- de Almeida Araújo IM, da Silva RM, Bonfim IM, Fernandes AF. Nursing communication in nursing care to mastectomized women: a grounded theory study. Rev Lat Am Enfermagem. 2010;18(1):54-60. doi: 10.1590/s0104-11692010000100009.
- Meleis Al. Theoretical Nursing: Development and Progress. Lippincott Williams & Wilkins; 2011.
- Ross JG, Bruderle E, Meakim CH, Willens JS, Holmwood JA. Development of formative capstone simulations to prepare novice students for initial clinical practicum. J Nurs Educ. 2016;55(10):587-9. doi: 10.3928/01484834-20160914-09.
- 22. Avan H, Koç ET, Vural B. Therapeutic communication and play with child. Kırşehir Ahi Evran Üniversitesi Sağlık Bilimleri Dergisi. 2020;1(2):62-9.
- 23. Bussard ME, Lawrence N. Role modeling to teach communication and professionalism in prelicensure nursing students. Teach Learn Nurs. 2019;14(3):219-23. doi: 10.1016/j.teln.2019.04.005.
- 24. Uhm JY, Ko Y, Kim S. Implementation of an SBAR communication program based on experiential learning theory in a pediatric nursing practicum: a quasi-experimental study. Nurse Educ Today. 2019;80:78-84. doi: 10.1016/j.

- nedt.2019.05.034.
- Kudubes AA, Ayar D, Zengin H. Therapeutic communication skills level among students undertaking the pediatric nursing course and the associated influencing factors. J Pediatr Nurs. 2023;73:34-43. doi: 10.1016/j.pedn.2023.08.015.
- Loghmani L, Borhani F, Abbaszadeh A. Factors affecting the nurse-patients' family communication in intensive care unit of Kerman: a qualitative study. J Caring Sci. 2014;3(1):67-82. doi: 10.5681/jcs.2014.008.
- 27. Ghorbani F, Zarei E, Mohamadi K. The relations of family members in the post-corona period based on the goals of developing a successful and healthy family in the horizon of 1410 and according to Setir's point of view. Mil Psychol. 2024;15(3):11-39. [Persian].
- Pourbandbani MA, Sadeghi R, Salsali M, Borhani F. Effective factors on active participation of clinical nurses to solution of ethical issues: master nursing students' perspective. Nurs Midwifery J. 2013;11(1):42-50. [Persian].
- Han S, Yoo J, Kang K. Development and validation of the therapeutic communication scale in nursing students. Healthcare (Basel). 2024;12(3):394. doi: 10.3390/ healthcare12030394.
- 30. Gholami S, Ahmadi S, Hormati H. Designing a curriculum model (teaching) empathy skills in the secondary schools and evaluating its effectiveness. Journal of Management and Planning in Educational System. 2021;14(1):311-40. doi: 10.52547/mpes.14.1.311. [Persian].
- 31. Mohammadi Nesbat P, Parchebafieh S, Fatah Moghadam L. The effectiveness of learning empathy skills on the responsibility of nursing students. Journal of Nurse and Physician Within War. 2023;11(38):28-35. [Persian].
- 32. Mutlu B, Azak M, Kalaycı S. Investigation of the relationship between clinical comfort-worry and self-efficacy perceptions of nursing students who have taken pediatric nursing course. OPUS International Journal of Society Researches. 2020;15(26):4293-315. doi: 10.26466/opus.659694.
- 33. Hojat M, Louis DZ, Markham FW, Wender R, Rabinowitz C, Gonnella JS. Physicians' empathy and clinical outcomes for diabetic patients. Acad Med. 2011;86(3):359-64. doi: 10.1097/ACM.0b013e3182086fe1.
- 34. Shariat SV, Kaykhavoni A. Empathy in medical residents at Iran University of Medical Sciences. Iran J Psychiatry Clin Psychol. 2010;16(3):248-56. [Persian].
- 35. Donovan LM, Mullen LK. Expanding nursing simulation programs with a standardized patient protocol on therapeutic communication. Nurse Educ Pract. 2019;38:126-31. doi: 10.1016/j.nepr.2019.05.015.
- 36. Managheb E, Bagheri S. The impact of empathy training workshops on empathic practice of family physicians of Jahrom University of Medical Sciences. Iran J Med Educ. 2013;13(2):114-22. [Persian].
- 37. Spiro H. What is empathy and can it be taught? Ann Intern Med. 1992;116(10):843-6. doi: 10.7326/0003-4819-116-10-843.